

Ramik Green®**KILLS RATS AND MICE****WEATHER RESISTANT RODENTICIDE FOR CONTROL OF MICE, RATS AND GROUND SQUIRRELS
INDOORS AND OUTDOORS****GUARANTEE:****Diphacinone**

(2-Diphenylacetyl-1, 3-Indandione).....0.005%

Registration No.: 11669

EPA Est. No. 61282-WI-1

PEST CONTROL PRODUCTS ACT

READ LABEL BEFORE USING**KEEP OUT OF REACH OF CHILDREN****WARNING**, contains the allergens Wheat, Soy and Fish
Contains Calcium Propionate at 1.0% as a preservative.**COMMERCIAL****FIRST AID:****If in eyes**, flush with water for 15 minutes.**If on skin**, wash with soap and water.**If swallowed**, give patient one or two glasses of water and induce vomiting by touching back of the throat with a finger or blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person.

Contact a physician or Poison Control Centre immediately.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Do not open pre-measured place packs. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.****ENVIRONMENTAL HAZARDS:** Keep out of lakes, streams, or ponds.

The use of an anticoagulant bait can present a significant hazard to predatory mammals and birds through secondary poisoning. Where possible remove and bury rodent bodies which are above ground.

FOR THE CONTROL OF: Mice and rats (indoors and outdoors) and ground squirrels, (Columbian, Franklin's, thirteen-lined and Richardson's) in farm yards, grain fields, nurseries, residential areas, turf (ornamental, sports) and vegetable gardens.**TOXICOLOGICAL INFORMATION:** Vitamin K₁ in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.**DISPOSAL:** Do not reuse empty container. Dispose of unused or spoiled bait in accordance with local requirements. Follow provincial instructions for any required cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For more information on the disposal of unused, unwanted product and the cleanup of spills, contact the **Provincial Regulatory Agency or the Manufacturer.****RODENT DISPOSAL:** Dispose of dead rodents in garbage or by burying.**NOTICE TO USER:** This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.**NOTICE TO BUYER:** Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product, and accepts the product on that condition.**DIRECTIONS FOR USE:** This product can be used in wet or damp locations such as river banks, railroad tracks, loading areas; in or around rat holes, under granaries, along fences, garbage dumps, etc.

All outdoor placements must be made in tamper-proof bait boxes.

This product is equally effective in dry areas such as the insides of buildings and other protected areas.

Place baits in tamper-proof bait stations along walls, in corners, and in concealed locations where rats can feed undisturbed.

Bait placements should be checked daily to make sure that an uninterrupted supply of bait is available. Maintain an uninterrupted supply of bait for at least 15 days. Continue baiting until all signs of feeding have stopped. Replace damp and mouldy bait.

RATS: Place bait 6 to 9 metres apart. At least 115 grams of bait should be used per location.**MICE:** Place bait 2 to 3 metres apart. At least 115 grams of bait should be used per location. Bait placements should be kept permanently to prevent new infestation.

GROUND SQUIRRELS:

Hand Baiting: Place 15 grams (200 pellets) of bait in each hole. Revisit the site after 48 hours and rebait all holes where the bait has been consumed or removed. Do not rebait holes where the bait has been undisturbed. A third visit may be necessary for full control.

Bait Station Method: Determine the approximate population of ground squirrels by a visual count over a 2 day period. Establish protected bait stations or tamper-proof bait boxes every 30 to 60 metres. Use one box for every 50 animals counted. Set 500 grams of bait per station. Check daily and maintain an uninterrupted supply for 3 weeks or until feeding has stopped.

IMPORTANT:

For best results, keep this product in a closed container, free from odours which may contaminate the bait and reduce acceptability. Rodents may feed several times before the effects are noticed. Several days may be required for death to occur.

Medical **1-800-498-5743**

? **1-800-621-8829**

Manufactured By:
Hacco, Inc.
110 Hopkins Drive
Randolph, WI 53956

Code (NG.NI) NA.NI

Item No.: 112952
Net Contents: 10 kg
UPC: 0 23626 01160 (1)
Format: UPC-A

Ramik® is a Registered Trademark of HACCO, Inc.