Ramik® Mouse Maze™

RODENTICIDE
KILLS MICE

GUARANTEE:
Diphacinone
(2-Diphenylacetyl-1,3-Indandione).....0.005%

Registration No.: 23739 EPA Est. No. 61282-WI-1
PEST CONTROL PRODUCTS ACT

READ LABEL BEFORE USING

DOMESTIC

KEEP OUT OF REACH OF CHILDREN

DIRECTIONS: Place the RAMIK® MOUSE MAZE™ in locations where mice are most likely to find and consume the bait. Place the bait in concealed corners, along walls, where mice feed. Place the bait at 1.5 to 3m intervals where mice are active. Maintain an uninterrupted supply for a period of 15 days or more.

USE LIMITATIONS: Bait MUST either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in contact with food.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Rubber gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin and clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS. Keep out of lakes, streams or ponds.

FIRST AID:
If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If swallowed, give patient 1-2 glasses of water and induce vomiting by inserting finger at the back of the throat, repeat until fluid is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: If swallowed by humans, pets, wildlife or domestic animals, this material may reduce the clotting ability of the blood and cause bleeding. Vitamin K_1 in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Dispose of empty container in household garbage. Dispose of unused bait in accordance with local requirements.

RODENT DISPOSAL: Dispose of dead rodents in garbage or by burying.

Medical **1-800-498-5743**
? **1-800-621-8829**

Manufactured By:
Hacco, Inc.
110 Hopkins Drive
Randolph, WI 53956

Code (NG.NT) NA.NI

Item No.:
Net Contents:
UPC:
Format:

This specimen label is intended for use only as a guide in providing general information regarding the directions, warning and cautions associated with the use of this product. As with any rodenticide, always follow the label instructions on the package before using.