

Havoc[®] Rodenticide Pellets

KILLS RATS AND MICE

For Control of Commensal Rats and House Mice

Second Generation Anticoagulant

Kills Warfarin-Resistant Norway Rats and House Mice. Rodents may consume a lethal dose in one feeding, with the first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT:

Brodifacoum -

3-[3-(4'-bromo-[1,1'-biphenyl]-4-yl)-
1,2,3,4-tetrahydro-1-naphthalenyl]-4-
hydroxy-2H-1-benzopyran-2-one0.005%

OTHER INGREDIENTS¹:.....99.995%

TOTAL100.000%

¹ Contains Bitrex[®] brand denatonium benzoate

EPA Reg. No. 100-1052-61282

EPA Est. No. 61282-WI-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If swallowed

- Call a poison control center, doctor, or **HOT LINE NUMBER** immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal)

Call **1-800-498-5743**

Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)

Call **1-800-424-9300**

NOTE TO PHYSICIAN OR VETERINARIAN: This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administrations of vitamin K₁ are indicated, as in poisoning from overdose of bishydroxy coumarin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (**not** mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to 2 weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. Keep away from humans, domestic animals and pets. Wash hands after handling bait.

IF BAIT IS EATEN BY ANIMALS OR PETS, CALL A VETERINARIAN AT ONCE.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under 6 years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway rats, Roof rats and House mice in and around homes, industrial, commercial, agricultural and public buildings, and similar manmade structures. This product may also be used in transport vehicles (ships, trains, aircraft) and in and around related port and terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS:

Norway and Roof Rats: Apply 4 – 16 oz. of bait (usually at intervals of 15-30 ft.) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

House Mice: Apply ¼ - ½ oz. of bait per placement. Space placements at intervals of 8-12 ft. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

Rats and Mice: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead rodents and left over bait properly. To prevent reinfestation, limit source of rodent food, water, and harborage as much as possible. Where a continuous source of infestation is present, maintain permanent bait stations and replenish with census or toxic bait as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store only in original container in a dry place inaccessible to children and pets.

Pesticide Disposal: Dispose of wastes resulting from the use of this product at an approved waste disposal facility or call your local solid waste agency for alternative disposal instructions. Never place unused product down any indoor or outdoor drain.

Pesticide Container: 1. Outer Bait Container: Do not reuse empty container except for holding additional Havoc® Rodenticide Pellets. Dispose of in a sanitary landfill or by other procedures if allowed by state and local authorities.

2. Bait Pack: Do not reuse empty container. Place in trash.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

Manufacturer and Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Manufacturer, and Buyer and User assume the risk of any such use.

MANUFACTURER AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall Manufacturer or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF THE MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Manufacturer and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitations of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Manufacturer.

For non-emergency (e.g., current product information) call:
1-800-621-8829

Manufactured For:
Hacco, Inc.
110 Hopkins Drive
Randolph, WI 53956

Code (RR/NA) NU.NI

Item No.: 116373
Net Contents: 1 pound (454 g)
UPC: 095242 16380 (4)
Format: UPC-A

Havoc® is a trademark of a Syngenta Group Company
Bitrex® is a registered trademark of Macfarlan Smith Ltd.,
Edinburgh, Scotland.
Product of the United Kingdom
Formulated in the USA