

RESTRICTED USE PESTICIDE

DUE TO HAZARD TO NON-TARGET SPECIES

For retail sale to and use only by Certified applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING

(CONTAINS ADDITIONAL INFORMATION WHICH MAY NOT APPEAR ON PACKAGE LABEL.)

Prozap[®] Zinc Phosphide Oat Bait

FOR USE IN AIRFIELDS

For Use to Reduce Potential Air-Strikes with Birds Feeding on California Ground Squirrels.

KEEP OUT OF REACH OF CHILDREN

CAUTION

THIS LABELING, THE PACKAGE LABEL, AND THE BOOKLET FOR PROZAP[®] ZINC PHOSPHIDE OAT BAIT (EPA REG. NO. 61282-14) MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR PROZAP[®] ZINC PHOSPHIDE OAT BAIT BEFORE APPLYING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE PACKAGE LABEL AND ON THE BOOKLET AND ON THIS SUPPLEMENTAL LABEL MUST BE FOLLOWED.

EPA Reg. No. 61282-14 EPA Est. No. 61282-WI-1

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR USE IN AIRFIELDS

For Use to Reduce Potential Air-Strikes with Birds Feeding on California Ground Squirrels.

USE RESTRICTIONS:

This supplemental label authorizes this product to be used to control California ground squirrels (*Spermophilus beecheyi*) in bait station applications in secured infield areas at airfields.

PREBAITING: To increase acceptance of treated bait by California ground squirrels, prebait with two to three teaspoons of untreated oat groats in each bait station. Do not apply toxic bait until it becomes apparent that ground squirrels are readily visiting the bait stations and have consumed all or nearly all of the prebait.

APPLICATION DIRECTIONS: Secure bait stations at locations where they will be readily accessible to target ground squirrels, preferably near active burrows. Secure stations 20 to 100 feet apart, depending upon the density of the ground squirrel population. Use 1-2 teaspoons of bait per bait station.

Check bait stations 1 or 2 days after bait was applied. If no bait remains in a particular bait station, replenish it with an additional 1 – 2 teaspoons of bait. Do not replenish any bait station more than once unless there is clear evidence that ground squirrels are continuing to feed from it. While wearing gloves, collect and dispose of all dead animals and leftover bait properly.

Bait stations may be maintained in areas on permanent bases but should only be loaded with toxic bait when ground squirrels are active and accepting grain-based foods in numbers sufficient to promote effective control.

HACCO, INC.
110 Hopkins Drive
Randolph, WI 53956

Expires 07/31/2011